



Delstrigo

nam aidsmap

What is *Delstrigo*?

Delstrigo is a medication used to treat HIV. It is a combination of three separate antiretroviral drugs in one pill, taken once a day.

It combines 100mg of doravirine, 300mg of lamivudine and 245mg of tenofovir disoproxil in a yellow tablet with '776' on one side of the tablet.

How does *Delstrigo* work?

Delstrigo combines three antiretroviral drugs in one pill. Two of the drugs (lamivudine and tenofovir disoproxil) are from a class of drugs known as NRTIs (nucleoside/nucleotide reverse transcriptase inhibitors). The third drug, doravirine, is a non-nucleoside reverse transcriptase inhibitor. Each drug class works against HIV in a different way.

The aim of HIV treatment is to reduce the level of HIV (the 'viral load') in your body until it is undetectable – usually less than 50 copies of virus per ml of blood. Taking HIV treatment and having an undetectable viral load protects your immune system and stops HIV being passed on to someone else during sex.

How do I take *Delstrigo*?

You should take *Delstrigo* once a day. *Delstrigo* can be taken with or without food.

HIV treatment works best if you take it every day, ideally at the same time each day. It may help to set an alarm, e.g. on your mobile phone, to remind you. If you forget to take a dose of *Delstrigo* and realise within 12 hours of the time you usually take it, take it as soon as possible, then take your next dose at your usual time. If you realise more than 12 hours late, don't take a double dose, just skip the dose you've forgotten and then carry on with your normal routine.

What are the side-effects of *Delstrigo*?

All medicines have possible side-effects. It's a good idea to talk to a doctor, nurse or pharmacist about what to expect before you start taking any medication, and how to manage any side-effects which occur.

A full list of side-effects, including less common side-effects, can be found in the patient information leaflet that comes with *Delstrigo*.

Side-effects can be described as:

- Common – a side-effect that occurs in at least one in a hundred people (more than 1%) who take this drug.



- Uncommon – a side-effect that occurs in fewer than one in a hundred people (less than 1%) who take this drug.

Common side-effects of *Delstrigo* include: abnormal dreams, difficulty in sleeping, nightmare, depression, headache, dizziness, drowsiness, cough, an irritated or runny nose, nausea, diarrhoea, abdominal pain, vomiting, hair loss, rash, muscle disorders, fatigue, fever.

Does *Delstrigo* interact with other drugs?

You should always tell your doctor and pharmacist about any other drugs or medication you are taking. That includes anything prescribed by another doctor, medicines you have bought from a high-street chemist, herbal and alternative treatments, and recreational or party drugs.

Some medicines or drugs are not safe if taken together – the interaction could cause increased, dangerous levels, or it could stop one or both of the drugs from working. Other drug interactions are less dangerous but still need to be taken seriously. If levels of one drug are affected, you may need to change the dose you take. This must only be done on the advice of your HIV doctor.

You should not take *Delstrigo* if you are currently taking medicines from the following groups:

- products that contain St John's wort (a herbal remedy used for depression and anxiety)
- carbamazepine, oxcarbazepine, phenobarbital and phenytoin (used to treat epilepsy and prevent seizures)
- rifampicin and rifapentine (used to treat some bacterial infections such as tuberculosis)
- mitotane (a cancer treatment)
- enzalutamide (a treatment for prostate cancer)
- lumacaftor (treatment for cystic fibrosis)
- nafcillin (an antibiotic)
- telotristat ethyl (diarrhoea treatment)
- lesinurad (anti-gout treatment)
- bosentan (blood pressure treatment)
- dabrafenib (skin cancer treatment)
- modafinil (used to treat severe sleep disorders, chronic drowsiness).

Can I take *Delstrigo* in pregnancy?

If you are considering having a baby, or think you might be pregnant, talk to your doctor as soon as possible about which combination of anti-HIV medications would be right for you. It is important to take antiretroviral treatment during pregnancy to prevent passing HIV from mother to baby.



Delstrigo is not usually used by women who are pregnant or want to get pregnant as it has not been studied in women who are pregnant.

Talking to your doctor

If you have any concerns about your treatment or other aspects of your health, it's important to talk about these. For example, if you have any symptom or side-effect which may be from your treatment, or if you are finding it difficult to take your medication every day, one of your healthcare team will be able to help.

Building a relationship with a doctor may take time. You may feel very comfortable talking to your doctor, but some people find it more difficult, particularly when talking about sex, mental health, or symptoms they find embarrassing. It's also easy to forget things you wanted to talk about.

Preparing for an appointment can be very helpful. Take some time to think about what you are going to say. You might find it helpful to talk to someone else first, or to make some notes and bring them to your appointment. Our online tool *Talking points* may help you to prepare for your next appointment – **visit www.aidsmap.com/talking-points**

The editors have taken all reasonable care in the production of this publication. Neither NAM, nor the editors, can be held responsible for any inaccuracies or mis-statements of fact beyond their control. Inclusion of information on any treatment or therapy does not represent an endorsement of that treatment or therapy by NAM or the editors.

We recommend that this information should always be used in conjunction with professional medical advice.

This factsheet is produced by an organisation called NAM, and has been reviewed by members of our volunteer panels of people living with HIV and medical professionals. We welcome your feedback on our information resources.

NAM provides up-to-date and impartial HIV information. Please visit us at our website where you can read the latest HIV news and sign up for free email updates.

NAM, Acorn House, 314-320 Gray's Inn Road, London WC1X 8DP.

Email info@nam.org.uk

Web www.aidsmap.com

Published February 2019.

Due for review February 2022